

Cervical screening: the colposcopy examination



**Cervical
Screening**



Why am I going to the colposcopy clinic?

You've been asked to go for a further examination because your cervical screening has shown some cell changes on your cervix. These changes act as early warning signals that cervical cancer may develop in the future. It is important to remember that it is very rare for these changes to be cancer.



What is a colposcopy?

A colposcopy is a simple examination of the cervix using a colposcope (a type of magnifying glass). It lets the doctor or specialist nurse look more closely at the changes on your cervix to decide if you need treatment. The examination is similar to the screening test, so some women may find it a bit uncomfortable.

Colposcopy can be done safely during pregnancy and will not affect delivery of your baby, or your ability to get pregnant in the future.



Should I prepare in advance?

Some doctors and specialist nurses prefer not to do a colposcopy during your period. Your appointment letter will tell you how to rearrange your appointment, if necessary.

The examination takes only 15 minutes but you will need to allow up to an hour for your whole visit. You may bring a friend or relative with you.

For your own comfort, you may want to wear a loose skirt to your appointment. You will then have to remove only your underwear for the examination.

Some women have a slight discharge or bleeding after the examination. You may want to bring a sanitary pad just in case.



What happens before the examination?

First, the doctor or specialist nurse will want to ask you some questions. These will relate to your periods, the type of contraception you use, and any operations or illnesses you have had in the past. It is particularly helpful if you can remember the date of the first day of your last period before attending the clinic.



What happens during the examination?

If not wearing a skirt, you will be asked to undress from the waist down and to position yourself on a special couch which has padded supports for your legs. A nurse will be with you during the examination.

The doctor or specialist nurse will gently insert a speculum into your vagina, just like your initial screening test. Sometimes a repeat test will be carried out. Then the specialist will examine your cervix using the colposcope. This looks like a pair of binoculars on a stand and does not touch you or go inside you.

The specialist will dab liquids onto your cervix to highlight any areas of cell changes more clearly. A small sample of tissue (a biopsy) may be taken from the surface of the affected area. This may sting a little but it should not be painful.





What happens after the examination?

The specialist may be able to tell you immediately what they have found and what treatment, if any, is needed. You may be given another appointment to come back to discuss the results of any biopsy and any treatment you may need.

After the examination, you should feel well enough to continue with your usual routine. If you've had a biopsy, you may have a light bloodstained discharge for a few days but this is nothing to worry about. You should not have sex for up to five days to allow the biopsy site to heal. Do not use tampons until the discharge clears up.



What will the examination show?

The colposcopy and any biopsy results will show the type and extent of the cell changes on the cervix. This will show if you need treatment and, if so, what sort. Only very rarely will a biopsy show cell changes that have already developed into cancer. Surgery and other treatments are generally used to treat cervical cancer.



What if I need treatment?

Any treatment you may require is usually simple and can mostly be done under a local anaesthetic as an outpatient. The choice of treatment will depend on your particular case. Treatment may be carried out at your first visit, but in some cases you may have to come back to the clinic for your treatment.



What should I do after treatment?

After treatment, you may have a bloodstained discharge for six to eight weeks. During this time, when you have your period you need to use sanitary pads rather than tampons. It is also best to avoid heavy exercise and not to have sex. This will allow the cervix to heal as quickly as possible.



Will treatment affect my ability to have children?

Treatment of cervical cell changes will not affect your ability to get pregnant. However, some type of treatments may increase your risk of having a premature baby in the future. The doctor or specialist nurse will explain the risks of your treatment in detail.



Will I need to have check-ups?

Around six months after treatment you will be offered cervical screening again to check that your treatment has been successful. If your result is normal or shows mild or borderline changes, your sample will then be tested for human papillomavirus (HPV). This is known as the **test of cure**. If this test comes back negative for HPV, you will be at very low risk of ongoing cervical disease at that point in time, but it will be important that you attend for screening again in three years time.

If the cervical screening result shows moderate or severe changes, or if the HPV test comes back positive, you will be referred for another colposcopy.

Although one in five women will be invited back for another colposcopy, only a few will need treated again. This is because it can take longer than six months for your immune system to clear HPV after treatment.

Further information

If you have any further questions on your condition or treatment, do not hesitate to discuss these with your GP or colposcopy clinic. They will be happy to help you.

For more information, or this leaflet in another language or format, visit:

www.cancerscreening.hscni.net



Public Health Agency

12-22 Linenhall Street, Belfast BT2 8BS.

Tel: 0300 555 0114 (local rate).

www.publichealth.hscni.net

Find us on:



www.becancerawareni.info