Daily Fluid Balance & Prescription Chart Write in CAPITAL LETTERS or use addressograph **Special Instructions:** Hospital _____ First names: Consultant: Ward: Hospital no:_ DOB:_ Health and Care no:______Check_idenui For Hourly, Cumulative fluid recording today From their 16th birthday **FLUID INPUT (ml) FLUID OUTPUT (ml) INTRAVENOUS FLUID & MEDICINES* ORAL FLUID** URINE **BOWEL** *Include volume of infused medicines ORAL **ENTERAL** Hourly Grand **Balance** Total Total Amount Amount Fluid Type Fluid Type Fluid Type Fluid Type Fluid Type Fluid Type OUT OUT Amount Total Amount 08.00 09.00 10.00 11.00 12.00 13.00 14.00 15.00 16.00 17.00 18.00 19.00 20.00 21.00 22.00 23.00 24.00 01.00 02.00 03.00 04.00 05.00 06.00 07.00 INTAKE 8am Day 8pm Night 8am Total OUTPUT 8am Day 8pm Night 8am Total Urine Oral Bowel Enteral Other 24 hour Fluid Balance (ml) Intravenous Other **Balance Grand Total IN**

Grand Total OUT

Write in CAPITAL LETTERS	or use addressograph
Surname:	
First names:	
Consultant:	Ward:
Hospital no:	DOB:
Health and Care no:	Check Ideliury

Recent Weight	_ kg
Weighed/ / Estimated	

Yesterday's Date	Grand total in	Grand total out	Balance			

	Composition of common intravenous solutions (mmol/L)						
Solution	[Na ⁺]	[CI ⁻]	[Glucose]	[Lactate]	[K ⁺]		
5% Glucose (Dextrose)	0	0	278	-	-		
0.45% Sodium Chloride	77	77	0	-	-		
0.45% Sodium Chloride + 5% Glucose (Dextrose)	77	77	278	-	-		
0.9% Sodium Chloride	154	154	0	-	-		
Compound Sodium Lactate - Hartmann's solution	131	111	0	29	5		
Plasma substitutes: gelatins, etherified starches	140 - 154	118 - 154	0	0 - 30	0 - 5		
Plasma	135 - 145	95 - 108	3.5 - 7.0	0.4 - 2.2	3.5 - 5.0		

FLUID PRESCRIPTION ADVICE FOR ADULTS*

Fluid therapy should involve the consideration of:-

- **RESUSCITATION** = **B** Fluid bolus volume for hypovolaemic shock.
- ROUTINE MAINTENANCE = M Varies with clinical state.
- REPLACEMENT Correction of any obvious fluid deficit = D and ongoing losses = o (e.g. vomiting, drainage, insensible, diarrhoea).

- **REDISTRIBUTION** Patients with sepsis, severe renal, liver or cardiac disease, malnutrition, refeeding issues or post-operative.
- REASSESSMENT include clinical assessments & glucose and U&E at least 24 hourly.
- Use the **oral** or **enteral** route whenever possible for giving fluids.
- Daily insensible loss = 500 800 ml.

Types of Fluid

Sodium chloride 0.9% provides the most important extracellular ions. It is indicated when RESUSCITATION by a fluid bolus is needed for shock and in sodium depletion. The administration of large volumes may give rise to sodium accumulation, oedema, and hyperchloraemic acidosis. Compound sodium lactate (Hartmann's solution: contains 5 mmol/L of potassium) can be used instead of isotonic sodium chloride solution during or after surgery, or in the initial management of the injured or wounded; it may reduce the risk of hyperchloraemic acidosis. 5% glucose (dextrose) is an important source of free water for maintenance, but should be used with caution as excessive amounts may cause dangerous hyponatraemia, especially in the elderly.

ROUTINE MAINTENANCE fluids replace the normal fluid content of oral food intake, insensible loss & urinary output and are prescribed to provide optimal hydration in patients unable to fully use the oral or enteral route. Maintenance fluids should take into account the volume of fluid to deliver IV medications (antibiotics, analgesics). The total (oral, IV drugs and prescribed fluids) volume prescribed in healthy adults (without excess fluid losses) should be of the order of 30 ml/kg/day up to a maximum of 2.5L. Consider prescribing less fluid (e.g. 20-25 ml/kg/day) for patients who are older, frail, have renal impairment or cardiac failure. Consider using ideal body weight for obese patients.

Sodium requirement - I to 2 mmol/kg/day, so it is rarely necessary to give more than I litre of sodium chloride 0.9% or Hartmann's solution per day for maintenance IV fluids. Antibiotic and analgesic infusions may already provide some of this. Potassium requirement - I mmol/kg/day, adjusted according to the serum potassium. Phosphate, Magnesium - monitor & correct.

Many patients have specific needs to cover REPLACEMENT and/or REDISTRIBUTION of fluid and electrolytes.

REPLACEMENT of deficits & on-going losses - prescription should reflect the electrolyte composition of the fluid being lost. Gastric losses - replace volume for volume with sodium chloride 0.9% with added potassium as required. Lower Gastrointestinal losses - replace with Hartmann's solution or, if extra potassium is needed, sodium chloride 0.9% with

Some patients have problems of internal REDISTRIBUTION and may develop sodium and water excess (leading to oedema and weight gain) which frequently can occur in the context of a low intravascular volume (and associated low urine output). Prescribing appropriate IV fluids for patients with redistributive type problems is particularly difficult since too little leads to intravascular hypovolaemia, low blood pressure, poor urine output and poor tissue perfusion, whilst too much may promote more oedema. In these patient groups, formulae-based equations should be used with caution. Fluid restriction may be needed and should be guided by senior input and regular REASSESSMENT.

Senior help should always be sought for complex on-going fluid losses, when the balance between fluid overload and deficit is unclear, for complex redisribution issues and especially when patients have diminished organ reserve.

*Based on NICE CG174

Indica	Indications - all that apply: Fluid Bolus volume, Deficit, On-going loss volume, Maintenance, Drug Prescription				* Medicines must be r	* Medicines must be recorded in Drug Kardex			** Model name, Serial number.					
\	Date	Time	Volume	Infusion Fluid/Type	Additives *	Rate ml/hour Range	Prescriber's Signature	Administered By	Checked By	Batch/Lot No. & Expiry Date	Pump Details **	Start Time	Finish Time	Volume given
				a										
				b										
				©										
				a										
				e										
				(f)										
				(g)										
				(b)										
				<u>(i)</u>				1						
				①										