

Smoking damages your eyesight

When you smoke, you are at increased risk of blindness

Smoking can:

- double your risk of cataracts
- damage vital blood vessels in your eyes
- make you 4 times more likely to develop macular degeneration

Stopping smoking at any age can reduce your risk. It's never too late to stop and there are over 600 free stop smoking services in Northern Ireland that can help. Look for the stop smoking services logo or for more information and to find your nearest service, visit www.stopsmokingni.info

HSC Health and Social Care Produced by the Public Health Agency
www.publichealth.hscni.net

03/19

Smoking and sight loss (leaflet and poster)

These resources outline the link between smoking and sight loss, highlighting how smoking significantly increases the risk of conditions such as cataracts, macular degeneration and retinopathy. They also signpost smokers to the free stop smoking services that are available across Northern Ireland to help them quit.

Published

14 March 2019

Last reviewed

27 March 2019

Details

Format

2 page A4 leaflet, A4 poster - PDF only

Target group

Smokers

Downloads

Attachment	Size
Smoking and sight loss leaflet 03.19 FINAL.pdf	929.04 KB
Smoking and sight loss poster 03.19 FINAL.pdf	542.43 KB

Tags

- [smoking](#)
- [sight](#)
- [smoking cessation](#)
- [cataracts](#)
- [macular degeneration](#)
- [retinopathy](#)

Share this page

- [Share by email](#)
- [Share on X](#)
- [Share on Facebook](#)
- [Share on LinkedIn](#)
- [Share on WhatsApp](#)
- [Download as PDF](#)

Was this page useful?

Yes No

Thank you for your feedback.

Sorry to hear that. [Tell us what went wrong.](#)