

	Chinese simplified
COVID-19 vaccination	冠状病毒疫苗接种
A guide for women who are pregnant or breastfeeding	孕口或哺乳期口女指南
Please read this before your vaccination	接种疫苗前，口先口口此内容
Find out more at www.nidirect.gov.uk/covid-vaccine	有关更多信息，口口口 www.nidirect.gov.uk/covid-vaccine
The COVID-19 vaccines available in the UK have been shown to be effective and to have a good safety profile. The current COVID-19 vaccines do not contain live coronavirus and cannot infect a pregnant woman or her unborn baby in the womb.	在英国可口得的冠状病毒疫苗已被口明是有效的，并且具有良好的安全性。当前的冠状病毒疫苗不含活病毒，也无法感染孕口或子口内未出生的口儿。
COVID-19 vaccination in pregnancy	孕口接种冠状病毒疫苗
Although clinical trials on the use of COVID-19 vaccines during pregnancy are not advanced, the available data do not indicate any harm to pregnancy.	尽管尚未口行有关在口孕期口使用冠状病毒疫苗的口床口口，但口有数据并未表明口口孕有任何危害。
In addition to any pre-existing risk factors a pregnant woman may have, in the later stages of pregnancy women are at increased risk of becoming seriously unwell with COVID-19. If this happens, your baby is two to three times more likely to be born prematurely, which can affect their long-term health.	除了孕口可能存在的任何口先存在的危口因素外，在孕口的后期口段，口女患了冠状病毒而造成口重不适的口口也越来越高。如果口生口种情况，您的口儿早口的可能性会高出两到三倍，口可能会影响他口的口期健康。
The Joint Committee on Vaccination and Immunisation (JCVI) has therefore advised that pregnant women should be offered vaccination at the same time as the rest of the population, based on their age and clinical risk group.	因此，疫苗接种和免疫口合委口会（JCVI）建口，口根据孕口的年口和口床口口，口孕口和其他人群同口提供疫苗接种。
There is now extensive experience of the use of the Pfizer BioNTech and Moderna vaccines during pregnancy in the USA with no safety concerns. These vaccines are therefore the preferred vaccines to offer to pregnant women of any age.	口在，在美国已有广泛的口口在口孕期口使用口瑞BioNTech和Moderna疫苗，并且没有任何安全口患。因此，口些疫苗是提供口任何年口的孕口的首口疫苗。
Pregnant women who commenced vaccination with AstraZeneca, however, are advised to complete the course with the same vaccine.	但是，建口口开始使用AstraZeneca疫苗接种的孕口也要使用相同的疫苗来完成口口程。
What does this mean for me?	口口我意味着什么？
If you are pregnant you will be able to receive a COVID-19 vaccine at the same time as other women of your age:	如果您口孕了，您将可以与同口的其他女性同口接受冠状病毒疫苗：
<ul style="list-style-type: none"> You may wish to discuss having it with your doctor or midwife. 	<ul style="list-style-type: none"> 您可能希望与您的医生或助口士口口。
<ul style="list-style-type: none"> Pfizer and Moderna vaccines are the preferred vaccines for pregnant women of any age, because of more extensive experience of their use in pregnancy. 	<ul style="list-style-type: none"> 口瑞和Moderna疫苗是任何年口段孕口的首口疫苗，因口它口在妊娠中的使用口口更口丰富。
<ul style="list-style-type: none"> If you decide to have a COVID-19 	<ul style="list-style-type: none"> 如果您决定接种冠状病毒疫苗，口告

vaccine, please tell the vaccination team that you are pregnant so that this can be recorded.	知疫苗接种小口您已口孕，以便可以口口下来。
<ul style="list-style-type: none"> Pregnant women who commenced vaccination with AstraZeneca, however, are advised to complete the course with the same vaccine. 	<ul style="list-style-type: none"> 但是，建口开始使用AstraZeneca接种的孕口也要使用相同的疫苗来完成口口程。
If you have had the first dose of COVID-19 vaccine and then become pregnant or find out you are pregnant:	如果您已接种第一口冠状病毒疫苗，然后口孕或口口自己已口孕：
<ul style="list-style-type: none"> you may wish to discuss having the second dose with your doctor or midwife. 	<ul style="list-style-type: none"> 您可能想要与您的医生或助口士口口第二口的使用。
<ul style="list-style-type: none"> unless you had serious side effects, you are advised to complete the course with the same vaccine during your pregnancy or you can wait until you have had your baby. 	<ul style="list-style-type: none"> 除非您有口重的副作用，否口建口您在口孕期口使用相同的疫苗完成整个口程，或者可以等到您的孩子出生后再口行。
Breastfeeding	哺乳
There is no known risk associated with any current COVID-19 vaccines whilst breastfeeding. The JCVI advises that breastfeeding women may be offered any suitable COVID-19 vaccine.	母乳喂养期口，目前没有任何已知的冠状病毒疫苗相关的口口。JCVI建口可以口母乳喂养的口女提供任何合适的冠状病毒疫苗。
Talk to your doctor or midwife if you have any concerns.	如果您有任何疑口，口咨口您的医生或助口士。
There is no need to avoid pregnancy after COVID-19 vaccination.	接种冠状病毒疫苗后无需避孕。
The Royal College of Obstetricians and Gynaecologists (RCOG) and Royal College of Midwives (RCM) have a decision guide and other useful information on COVID-19 vaccines and pregnancy (www.rcog.org.uk and www.rcm.org.uk/guidance-for-pregnant-women)	皇家口口科学院（RCOG）和皇家助口士学院（RCM）口有关于冠状病毒疫苗和妊娠的决策指南和其他有用信息(www.rcog.org.uk and www.rcm.org.uk/guidance-for-pregnant-women)
Vaccination, helping to protect those most vulnerable.	接种疫苗，有助于保口最脆弱的人群。
Information correct at time of publication. For the latest version of this factsheet, visit the PHA website www.publichealth.hscni.net/publications	出版口信息正确。有关此情况口明口的最新版本，口口口PHA网站 www.publichealth.hscni.net/publications
© Crown Copyright 2020. This information was originally developed by Public Health England and is used under the Open Government Licence v3.0	© Crown Copyright 2020.此信息最初由英国公共口生部口开口，并在“开放政府口可口”v3.0下使用
Public Health Agency	公共口生署
Tel: 0300 555 0114 (local rate)	口口:0300 555 0114 (当地口口口率)
Find us on:	在此找到我口：