

Chinese Simplified translation of *COVID-19 vaccination - Your guide to booster vaccination* 12/21

English	Chinese Simplified
COVID-19 vaccine	2019 冠状病毒病（COVID-19）疫苗
Your guide to booster vaccination	您的疫苗加强剂指南
People aged 18 years and over, health and social care workers and those aged 16 and 17 who are at risk are being offered a booster dose of coronavirus (COVID-19) vaccine.	目前正在向 18 岁及以上的人、卫生和社会护理工作者以及存在风险的 16 岁和 17 岁的人提供冠状病毒（COVID-19）疫苗加强剂。
What is coronavirus or COVID-19?	什么是冠状病毒或 COVID-19?
COVID-19 is a very infectious respiratory disease caused by the SARS-CoV-2 virus and is more serious in people over 40 years of age and those with certain health conditions.	COVID-19 是一种传染性很强的呼吸道疾病，由 SARS-CoV-2 病毒引起，在 40 岁以上的人和具有某些健康问题的人中更为严重。
Why are you being offered a COVID-19 booster?	为什么向您提供 COVID-19 疫苗加强剂?
Like some other vaccines, levels of protection may begin to wane over time. The booster will help to reduce the risk of you needing admission to hospital due to COVID-19 infection this winter. Protection against severe disease from the first 2 doses seems to decline very slowly. Boosting your immunity should help to extend your protection into next year and give you longer term protection.	像其他一些疫苗一样，其保护水平可能会随着时间的推移开始减弱。疫苗加强剂将有助于降低您在今年冬天因 COVID-19 感染而需要住院的风险。从前 2 剂得到的对严重疾病的保护似乎在非常缓慢地下降。提高您的免疫力应该有助于将您的保护延长到明年，并可能对新变种提供更广泛的保护。

Chinese Simplified translation of *COVID-19 vaccination - Your guide to booster vaccination 12/21*

When should you have a booster?	您应该什么时候接种疫苗加强剂？
The booster is being offered to those most vulnerable first and will then be rolled down to younger age groups. Your appointment should be at least 3 months from your last dose, but this may be later depending on your age group or level of risk. For more on this, visit www.nidirect.gov.uk/covid-vaccine	正在首先向那些最易受感染的人提供加强剂，然后将推广至更年轻的年龄组。您的预约应该在您最后一次接种后至少 3 个月，但可能会根据您的年龄组或风险水平而更迟联系您。欲了解更多相关信息，请访问： www.nidirect.gov.uk/covid-vaccine 。
Which vaccine will you be offered?	您将被提供哪种疫苗？
You will be given a booster dose of either Pfizer or Moderna vaccine. Both vaccines boost well and have already been given to millions of people in the UK. Studies have shown that you only need a half dose of Moderna to boost the immune system well. This half dose of Moderna is expected to have low rate of side effects including myocarditis. See page 3 for more information.	您将获得辉瑞（Pfizer）或莫德纳（Moderna）的加强疫苗。这两种疫苗都能有效加强免疫力，并且已在英国提供给数百万人使用。研究表明，您只需要半剂量的莫德纳疫苗即可很好地增强免疫系统。预计这半剂量的莫德纳疫苗的副作用（包括心肌炎）发生率很低。欲了解更多信息，请参见第 3 页。
You will be offered the right vaccine for you which may be the same or different from the vaccines that you had before.	您将获得适合您的疫苗，该疫苗可能与您之前接种的疫苗相同或不同。

Chinese Simplified translation of *COVID-19 vaccination - Your guide to booster vaccination 12/21*

Common side effects	常见的副作用
As with your previous dose the common side effects are the same for all COVID-19 vaccines used in the UK, and include:	与您之前接种的疫苗一样，在英国使用的所有 COVID-19 疫苗的常见副作用都相同，包括：
<ul style="list-style-type: none">• having a painful, heavy feeling and tenderness in the arm where you had your injection. This tends to be worst around 1 to 2 days after the vaccine	<ul style="list-style-type: none">• 注射部位的手臂有疼痛，沉重的感觉和压痛。在疫苗接种后约 1-2 天，这种情况往往最严重。
<ul style="list-style-type: none">• feeling tired	<ul style="list-style-type: none">• 感觉累
<ul style="list-style-type: none">• headache	<ul style="list-style-type: none">• 头痛
<ul style="list-style-type: none">• general aches, or mild flu like symptoms	<ul style="list-style-type: none">• 全身痛，或轻度流感样症状
You can rest and take paracetamol (follow the dose advice in the packaging) to help make you feel better. Although feeling feverish is not uncommon for 2 to 3 days, a high temperature is unusual and may indicate you have COVID-19 or another infection.	您可以休息，并服用扑热息痛（paracetamol）（按照包装上的剂量建议），以使您感觉好些。尽管在两到三天内感觉发烧并不罕见，但很少会出现高烧，这可能表明您感染了冠状病毒或有其他感染。
Although a fever can occur within a day or 2 of vaccination, if you have any other COVID-19 symptoms or your fever lasts longer, stay at home and arrange to have a test. Symptoms following vaccination normally last less than a week. If your symptoms seem to get	尽管在接种疫苗后一两天内可能会发烧，但是如果您有其他任何 COVID-19 症状或发烧持续时间更长，请留在家中并安排检查。接种疫苗后的症状通常持续不到一周。如果您的症状似乎变得更糟或是您担心，请致电您的全科医生（GP）。

Chinese Simplified translation of *COVID-19 vaccination - Your guide to booster vaccination* 12/21

worse or if you are concerned, contact your GP.	
You can also report suspected side effects of vaccines and medicines through the Yellow Card scheme. You can do this online by searching Coronavirus Yellow Card or by downloading the Yellow Card app.	您还可以通过黄卡计划（Yellow Card scheme）报告疑似的疫苗副作用和药物副作用。您可以通过搜索冠状病毒黄卡（Coronavirus Yellow Card）或下载黄卡应用程序（Yellow Card app）在线进行此操作。
If you had serious side effects after any previous dose you may be advised to avoid or delay further vaccination. You should discuss this with your doctor or specialist.	如果您在之前的任何接种后出现严重副作用，可能会建议您避免或推迟进一步接种疫苗。您应该与您的医生或专科医生讨论这个问题。
Serious side effects	严重的副作用
Worldwide, there have also been recent, very rare cases of inflammation of the heart called myocarditis or pericarditis reported after Pfizer and Moderna COVID-19 vaccines.	在全球范围内，最近出现了非常罕见的心脏炎症病例，称为心肌炎或心包炎，是在接种辉瑞或莫德纳 COVID-19 疫苗后被报告的。
These cases have been seen mostly in younger men within several days after vaccination. Most of these people recovered and felt better following rest and simple treatments.	这些病例主要见于接种疫苗后几天内的年轻男性。这些人中的大多数人在休息和简单治疗后感觉好转。
You should seek medical advice urgently if, after vaccination, you experience:	如果您接种完疫苗后有以下情况，应该立即寻求医疗建议：
<ul style="list-style-type: none"> • chest pain 	<ul style="list-style-type: none"> • 胸痛

Chinese Simplified translation of *COVID-19 vaccination - Your guide to booster vaccination 12/21*

<ul style="list-style-type: none"> • shortness of breath 	<ul style="list-style-type: none"> • 呼吸急促
<ul style="list-style-type: none"> • feelings of having a fast-beating, fluttering, or pounding heart 	<ul style="list-style-type: none"> • 感觉心跳节奏快，心颤动或心脏狂跳
Is there anyone who shouldn't have a booster?	有没有人不应该接种疫苗加强剂？
There are very few people who should not have a booster.	很少有人不应该接种疫苗加强剂。
If you have had a severe reaction to a previous dose of the vaccine you should discuss this with your doctor.	如果您对之前接种的疫苗有严重反应，您应该与您的医生讨论这个问题。
Can you still catch COVID-19 after having the vaccine?	您会从疫苗中感染冠状病毒吗？
The COVID-19 vaccination will reduce the chance of you suffering from COVID-19 disease. It may take a few days for your body to build up some protection from the booster.	已证明所使用的 COVID-19 疫苗可以减少您患 2019 冠状病毒病的机会。您的身体可能需要几天的时间才能建立起由疫苗得到的保护。
Like all medicines, no vaccine is completely effective – some people may still get COVID-19 despite having a vaccination, but this should be less severe.	像所有药物一样，没有一种疫苗是完全有效的——有些人尽管进行了疫苗接种，仍可能会感染冠状病毒，但这应该不那么严重了。
If you have not had the first vaccinations	如果您尚未接种之前的疫苗
If you have not yet had either of your first 2 doses of the vaccine you should have them as soon as possible.	如果您尚未接种前 2 剂疫苗中的任何一剂，则应尽快接种。

Chinese Simplified translation of *COVID-19 vaccination - Your guide to booster vaccination* 12/21

When can you have your booster if you've had a COVID-19 positive test result?	如果您的 COVID-19 结果呈阳性，什么时候可以接种加强剂？
You will still need the booster but you should wait at least 4 weeks from your COVID-19 infection.	您仍将需要加强剂，但您应该在感染 COVID-19 后至少等待 4 周。
Further information	额外信息
Visit coronavirus vaccination www.nidirect.gov.uk/covid-vaccine	访问冠状病毒疫苗信息 www.nidirect.gov.uk/covid-vaccine
See the PHA leaflet on what to expect after vaccination, which is available to download at www.pha.site/COVID19InfoMaterials	请参阅公共卫生署（PHA）有关接种疫苗后会发生什么的宣传单，该宣传单可在以下网址下载 www.pha.site/COVID19InfoMaterials
Please read the product information leaflet for more details on your vaccine, including possible side effects, by searching Coronavirus Yellow Card. You can also report suspected side effects on the same website or by downloading the Yellow Card app.	请通过搜索冠状病毒黄卡（Coronavirus Yellow Card），阅读产品信息手册，以了解有关疫苗的更多详细信息，包括可能的副作用。您还可以在同一网站上或通过下载黄卡应用程序（Yellow Card app）报告疑似的副作用。
https://coronavirus-yellowcard.mhra.gov.uk	https://coronavirus-yellowcard.mhra.gov.uk

Chinese Simplified translation of *COVID-19 vaccination - Your guide to booster vaccination 12/21*

You can read the information for UK recipients of the Pfizer and Moderna vaccines here:	您可以在此处阅读辉瑞和莫德纳疫苗的“给英国接受者的信息”：
https://coronavirus-yellowcard.mhra.gov.uk/productinformation	https://coronavirus-yellowcard.mhra.gov.uk/productinformation
Published September 2021 by the Public Health Agency.	由公共卫生署在 2021 年 9 月出版
© Crown copyright 2021. This information was originally developed by UK Health Security Agency and is used under the Open Government Licence v3.0	© 皇家版权 2021。此信息最初由英国卫生安全局（UK Health Security Agency）开发，并根据“第三版开放政府许可证”（Open Government Licence v3.0）使用。