Salmonella HSC Public Health Agency

What is Salmonella?

Salmonella are bacteria that cause a range of symptoms including diarrhoea, stomach cramps and sometimes vomiting and fever. It is one of the most common causes of food poisoning in the United Kingdom.

Who gets Salmonella infections?

Anyone can get *Salmonella* infection, but certain groups are more likely to get the infection:

- young children under 5 years old;
- travellers abroad.

How do you get infected with Salmonella?

Most people become infected by eating contaminated food. *Salmonella* bacteria can be present in undercooked poultry or meat, raw or undercooked eggs and unpasteurised milk.

It is impossible to tell from its appearance whether food is contaminated with *Salmonella*. It will look, smell and taste normal so correct handling and cooking are very important.

It is also possible to get *Salmonella* infection through contact with someone else who has the illness or contact with their contaminated items (for example when changing a nappy).

Rarer causes of infection include handling pet reptiles (including tortoises and terrapins) and exposure to contaminated water.

How can you avoid getting infected with Salmonella?

Wash your hands thoroughly with soap and warm water:

- before preparing and eating food;
- after handling raw food;
- after going to the toilet or changing a baby's nappy;
- after contact with pets and other animals;
- after working in the garden.

Avoid eating raw or lightly cooked eggs.

Do not drink untreated water from lakes, rivers or streams.

Avoid unpasteurised milk.

Do not keep reptiles or amphibians in households where there is a child under five years of age, or someone with an immune system that does not work properly (including people with cancer, AIDS or alcohol addiction).

Maintain high standards of hygiene when storing, handling or preparing food.

How can you maintain high standards of hygiene when storing, handling or preparing food?

Take chilled and frozen foods home quickly – then put into the refrigerator or freezer at once.

Keep cooked food away from raw food. Store raw foods below cooked or ready-to-eat foods in the fridge to prevent contamination. Cook food thoroughly, especially meat, so that it is piping hot. Follow cooking instructions on the pack.

If you reheat food, make sure it is piping hot all the way through. Only reheat food once.

Keep hot foods hot and cold foods cold. Don't just leave these foods sitting at room temperature.

Keep your refrigerator at 0–5° Centigrade. Get a thermometer to check temperatures inside the refrigerator.

Check 'use by' dates on food, and use foods within the recommended date on the pack.

Keep pets away from food, dishes and worktops.

Wash hands thoroughly before preparing and eating food, and after handling raw food

Keep all kitchen surfaces and equipment, including knives, chopping boards and dish cloths clean. Wash worktops and utensils between handling raw and cooked foods.

How can you prevent others in your household becoming infected with *Salmonella*?

Everyone in the house should wash their hands thoroughly with soap and warm water after going to the toilet and after handling soiled clothing or bedding, and before preparing/eating food.

Wash the infected person's laundry separately and on the hottest cycle of the washing machine for that particular fabric. Clean toilet seats, toilet bowls, flush handles, taps and wash hand basins after use with detergent and hot water, followed by a household disinfectant.

Everyone in the house should have their own towel and they should not use anyone else's.

The infected person should not prepare or serve food for others.

What are the symptoms of *Salmonella* and how long do they last?

Symptoms usually include diarrhoea and stomach cramps.

Other possible symptoms include nausea, vomiting, fever, headache, muscle aches and generally feeling unwell.

Symptoms usually develop 12–72 hours after exposure to the infection. Most cases start to resolve within three to seven days.

More severe illness can occur in young children, the elderly and people who have immune systems that are not working properly (including people with cancer, AIDS or alcohol addiction).

How are Salmonella infections treated?

There is no specific treatment for *Salmonella* infections. Most people will recover fully without treatment.

If you need advice on the treatment of your infection you should discuss this with your GP.

Always contact your GP if any of the following apply:

- your symptoms are severe or do not improve after a few days;
- you have a high temperature (fever) of 38°C (100.4°F) or over;
- you have symptoms of severe dehydration, such as sunken eyes and passing small quantities of dark, strong smelling urine;
- you have a baby with symptoms.
- you have an immune system that is not working properly (including people who have cancer, are having chemotherapy, or have HIV/ AIDS).

Do you need to stay off work or school?

Yes. While you are ill and have symptoms you are infectious to other people. You can return to work or school once you have been free from diarrhoea for 48 hours.

You should tell your employer you have had a *Salmonella* infection if you work with vulnerable groups such as the elderly, the young, those in poor health, or if you handle food.

Children should not attend school/childcare or play with other children until they are fully recovered and symptom free for 48 hours.



Public Health Agency, 12-22 Linenhall Street, Belfast BT2 8BS. Tel: 0300 555 0114 www.publichealth.hscni.net